



Schmidt, D. N. (2018). Determining climate change impacts on ecosystems: the role of palaeontology. *Palaeontology*, 61(1), 1-12. <https://doi.org/10.1111/pala.12335>

Peer reviewed version

License (if available):
CC BY-NC

Link to published version (if available):
[10.1111/pala.12335](https://doi.org/10.1111/pala.12335)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Wiley at <http://onlinelibrary.wiley.com/doi/10.1111/pala.12335/abstract> . Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available: <http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

CLIMATE CHANGE IMPACTS: THE ROLE OF PALAEOLOGY

Determining climate change impacts on ecosystems: the role of palaeontology

Climate change is projected to change the ecosystems on land and in the sea at rates that are unprecedented for millions of years. The fossil record is an archive of responses to climate change at a global ecosystem scale. While records are often fragmented, there are time intervals which – when thoroughly analysed with quantitative data – can provide valuable insights into the future of biodiversity on this planet. However, the most commonly used approach to derive projections of how ecosystems will look in the future are experiments on living organisms. By their nature, experiments are unlike the real world and cannot capture the ability of organisms to migrate, select, and evolve. They are often limited to a select few species and drivers of environmental change and hence cannot represent the complexity of interactions in ‘real’ ecosystems. This review provides an overview of projected impacts on marine ecosystems including: (1) the range of neontological methods, observations and their challenges; and (2) the complementary information that palaeontologists can contribute to this global challenge. I advocate that, in collaborations with other disciplines, we should aim for a strong visibility of our field and the knowledge it can provide for policy relevant assessments of the future.

Key words: Climate change, marine ecosystems, warming, acidification, PETM,

CLIMATE CHANGE IMPACTS: THE ROLE OF PALAEOONTOLOGY

Rising CO₂ levels are expected to radically alter our planet. Increasing atmospheric CO₂ radically alters the Earth system, with projected implications for ecosystems worldwide (e.g., Turley *et al.* 2010; Pörtner *et al.* 2014). The oceans, which cover 71% of the Earth's surface and represent >95% of its habitable environment, are likely to be strongly affected. The increase in atmospheric CO₂ alters the physical, chemical and biological properties of the ocean such as temperature, pH, stratification, and oxygen and nutrient availability (Pörtner, *et al.* 2014). Indirect consequences of a warmer world are increases in sea level and storminess (Stocker *et al.* 2013). Climate models project warming of the average surface ocean of 3°C, a loss of oxygen of the subsurface ocean of 3%, and a decrease in net primary productivity of 8% under business-as-usual scenarios by the end of the century (Lefort *et al.* 2015). High resolution Earth system models predict that the rise in CO₂ will result in carbonate undersaturation in high latitude surface waters, with respect to aragonite, by the middle of the century (Orr *et al.* 2005). By 2050, using the same scenario, physical drivers of biology, physiology, and ecology in most of the ocean will be outside their background variability (Henson *et al.* 2017).

However, what is much less well understood than the physical changes, is their impact on biodiversity and benefits gained by humans from ecosystems, termed 'ecosystem services' (e.g., Turley, *et al.* 2010; Pörtner, *et al.* 2014). Ecosystem services include (1) provisioning services such as food; (2) regulating services that affect climate; (3) cultural services that provide recreational, aesthetic, and spiritual benefits; and (4) supporting services such as photosynthesis, and nutrient cycling (Millennium Ecosystem Assessment 2005). Amongst the most pressing questions concern physiological impacts on individuals (Pörtner, *et al.* 2014), how species change their

CLIMATE CHANGE IMPACTS: THE ROLE OF PALAEOLOGY

distribution, abundance and phenology (Edwards *et al.* 2001), if the consequences of these changes in fundamental ecosystem structure impact regulatory and provisioning services (Poloczanska *et al.* 2013), and whether or not species can use their natural plasticity or evolve (Collins *et al.* 2014) to dampen the impact of climate change (Lohbeck *et al.* 2012; Collins and Bell 2004). Quantifying the full range of natural responses to climate change remains a key challenge.

Our current knowledge is based on several lines of evidence. Responses to ongoing climate change show shifts in abundance, geographical distribution, and migration patterns of marine organisms (Edwards, *et al.* 2001; Beaugrand *et al.* 2002; Beaugrand and Reid 2003). Poleward displacement of plankton, making use of the new opportunities as warmer waters expand, has been as fast as 100s of km/decade (Poloczanska, *et al.* 2013). These species specific changes in range expansion result in changes in food web structures (Kortsch *et al.* 2015) thereby impacting fisheries (Cheung *et al.* 2013).

These observations are supported by model projections of significant changes in the distribution, biomass and maximum size of pelagic communities by the year 2100 (Lefort, *et al.* 2015). Biomass and maximum size is projected to increase in the high latitudes but decrease in low and mid latitudes, impacting on large species with high metabolic needs because of reductions in food availability (Lefort, *et al.* 2015). It has been found that 14% of the global ocean experienced important species turnover (defined as the sum of species invasion and extirpation). In this analysis, while 6% - 44% of the ocean, mainly in the tropics, is expected to experience reduced biodiversity (Beaugrand *et al.* 2015) biodiversity is predicted to increase in high latitudes.

CLIMATE CHANGE IMPACTS: THE ROLE OF PALAEOONTOLOGY

Observations in the field have been corroborated by the results of experiments in laboratories and mesocosms that help provide mechanistic understanding of the underlying causes (Pörtner, *et al.* 2014). These experiments revealed differential vulnerability across the life stages of sea urchins (Dupont *et al.* 2012), changes in energy allocation to physiological processes (e.g. muscle wastage in brittle stars at continued calcification (Wood *et al.* 2008)) and the importance of energy availability in resilience to climate change in mussels (Thomsen *et al.* 2010), to name but a few. These experiments highlight that not all species or processes have the same vulnerability and some are not impacted at all (Wall *et al.* 2015).

By their nature, short-term experiments lack realism and cannot capture the organism's ability to migrate, to select from a vast population size, and ultimately the evolution of the species. The experiments need to focus on short-lived organisms that deal well with the laboratory conditions. Adaptive evolution is more likely to occur in species with fast generation times and large population sizes, such as marine microbes (Krueger-Hadfield *et al.* 2014; Scheinin *et al.* 2015). Long-term, multigenerational experiments are rare (Lohbeck, *et al.* 2012; Collins and Bell 2004) and often limited to asexual offspring of species with extremely short generation times under simplified environmental conditions. The results of these experiments suggest that adaptive evolution increases resilience, thereby potentially reducing the size of the impact on the ecosystem (Collins, *et al.* 2014).

A complementary approach to assess impacts of climate change and to generate such mechanistic understanding is based on the fossil record, wherein climate change

CLIMATE CHANGE IMPACTS: THE ROLE OF PALAEOONTOLOGY

events occurred at different rates and amplitudes (Hönisch *et al.* 2012). Therefore, the fossil record is an ideal test bed for model projections of changes of distribution, size, abundance and biomass. Marine sediments provide archives of biotic and climate variability in response to abrupt climate change that are recorded with high temporal resolution, good preservation and global distribution. They preserve a wealth of information including abundance, size, biomineralization, ontogeny and reproductive strategy (in some species), and migration and extinction. The ability to compare the open ocean with the coast, the surface with the deep ocean, the land and the ocean for a vast number of species holds incredible potential for the contribution of palaeontology to this ongoing debate. Exploiting the fossil record of environmental change and its biotic response will allow the rules governing biotic responses to climate change in the past to be identified, thus informing predictive models of the future.

Here, I will assess how past records of biotic response to climate change can contribute to quantifying impacts of future climate change in the next decades. I will start with the independent environmental framework necessary for such assessment and discuss the impacts of climate change on biogeography and calcification, with a focus on the marine microfossil record. Using the Paleocene Eocene Thermal Maximum (PETM) as a case study, I will highlight the power of an integrated analysis of climate data, modelling, and marine and terrestrial fossil records.

ENVIRONMENTAL FRAMEWORK

Interrogating the fossil record of climate change critically hinges on the accuracy of the reconstruction of the physical environment. Too often, circumstantial evidence is

CLIMATE CHANGE IMPACTS: THE ROLE OF PALAEOONTOLOGY

used to derive the physical change without independent information. Over recent years many proxies have appeared that allow us to quantify, within their constraints, environmental changes (e.g., Henderson 2002; Hönisch, *et al.* 2012). There are two main approaches to reconstructing the environment: (1) climate proxy compilations based on biotic and abiotic information stored in the fossil record (Robinson *et al.* 2008; Dunkley Jones *et al.* 2013); or (2) climate models (Lunt *et al.* 2010b; Stocker, *et al.* 2013); and, ideally, there would be a combination of both (Lunt *et al.* 2010a; Tindall *et al.* 2010). For climate models, the desired geographical resolution (Bopp *et al.* 2013; Dutkiewicz *et al.* 2015) has to be balanced against the ability to run these models for thousands of years (Ridgwell and Schmidt 2010). Without clear data or model output to attribute a biotic change to an environmental driver, circular reasoning is not uncommon. Ecological niche models (ENMs) provide the link between the physical climate models and distributions of species. ENMs estimate a species abiotic niche by correlating a known species occurrence with spatial environmental variables (Guisan and Zimmermann 2000; Pearson and Dawson 2003). They have been widely applied to modern distributions, mainly in the terrestrial record (Walther *et al.* 2002). They allow the quantification of changes in the abiotic niche in the past and then project these into the future (Stanton *et al.* 2014; Waterson *et al.* 2017). For example, Waterson *et al.* (2017) assessed if the environmental niches of several species of planktic foraminifera were stable between the last glacial maximum and today and if, therefore, species compositions can be used to reconstruct temperature difference using an ecological niche model. The strength of the fossil record makes it possible to distinguish between total range (presence absence) and the optimal niche (in this case high relative abundances/large body size (Schmidt *et al.* 2004; Schmidt *et al.* 2006). While the total niche of the investigated foraminiferal species did not change much

CLIMATE CHANGE IMPACTS: THE ROLE OF PALAEOLOGY

from the last glacial maximum to today, the optimal niche was more susceptible to environmental changes (Waterson, *et al.* 2017).

CLIMATE CHANGE IMPACTS ON MARINE MICROFOSSILS

Marine microfossils are an ideal system for investigating response to climate change: (1) they are very sensitive to a changing ocean environment (directly in contact with the changing environment, as opposed to metazoans) and cannot regulate their body temperature; (2) they have no known predator and, as such, are likely to react strongly to the abiotic environment with less important biotic interactions; (3) they have enormous population sizes and very few apparent barriers to gene flow, enabling a very rapid evolutionary response to selective pressures; (4) some groups preserve their entire ontogeny in their shell; and (5) they have an excellent global fossil record spanning 200 million years, and have survived periods of major climate change of different rates and amplitudes. Every sample contains thousands of specimens that also record environmental information via proxies thereby allowing direct links between environment and the biotic response.

The fossil record of marine microfossils spans multiple intervals of past rapid climate change over more than 100 million years (myrs). Today, marine microfossils have global distributions with clearly defined biogeographical zones suggesting strong environmental controls determined by a few dominant species. Their modern ecosystem functions range from primary production at the base of the food chain and the biological pump (phytoplankton) to carbonate and silicate production and therefore regulation of global biogeochemical cycles.

CLIMATE CHANGE IMPACTS: THE ROLE OF PALAEOLOGY

Biogeography changes

Changes in geographical ranges and abundance of microfossils have been applied to assess changes in climate in the past, most successful by CLIMAP (CLIMAP Project Members 1976) and its offspring the Margo project (Margo Project Members 2009). If combined with climate model that quantify the natural environment, these data can be used to interpret response to climate change in the past and project the future. We assume that plankton simply migrates without changes to their niche and this provides the foundation for using microfossil distributions in transfer functions for environmental reconstructions. Niche models, though, challenge the assumption of niche stability in foraminifers over the last 20 thousand years (Waterson *et al.* 2016).

The dominant signal in marine plankton to temperature changes is habitat tracking, i.e. an expansion of warm water species towards to pole in response to warming (Dowsett *et al.* 2012). Locally, this can lead to increases of warm water species in areas that previously did not support these species. For example, an annually resolved foraminiferal abundance record off the coast of California shows an increase in temperate/subpolar species over the last decades in response to the warming of the California Current (Field *et al.* 2006). Such high resolution records of response to climate change are rare but invaluable in assessing impact of environmental change.

CLIMATE CHANGE IMPACTS: THE ROLE OF PALAEOONTOLOGY

Combining the physiological understanding with climate models predicting future temperature, food and light suggests decrease of foraminiferal abundance and diversity in the tropics and subpolar regions and increase in the subtropics and around the poles for the future (Roy *et al.* 2015); though the model does not consider the impact of carbonate chemistry changes especially in high latitudes. Such assumptions are testable at various temperatures in the past across glacial interglacial cycles with warmer time intervals during the Eemian for example.

Calcification response – regulatory ecosystem services

A fundamental ecosystem function of foraminifers and coccolithophores is carbonate production, which regulates biogeochemical cycles, and is determined by abundance, size and weight of a population. Calcification is an energy demanding process and projected to be strongly affected by changes in CO₂ (Kroeker *et al.* 2010). Marine plankton in the modern ocean precipitate around 50% of the marine carbonate (Schiebel 2002) with the other half being produced on the shelf by a wide range of organisms including corals, calcifying algae, and molluscs. Over the course of the Cenozoic, pelagic carbonate production was dominated by coccolithophores, whilst today both coccolithophores and foraminifers contribute similarly to overall carbonate production.

Mass extinctions in the fossil record highlight the long duration of reduced carbonate accumulation (Henehan *et al.* 2016; Hull *et al.* 2011) and hence likely production, raising concerns about the future. Mechanistic understanding of the drivers of

CLIMATE CHANGE IMPACTS: THE ROLE OF PALAEOONTOLOGY

calcification is fundamental to forecasting impacts of climate change on calcification, the marine carbon cycle and feedback on the ocean CO₂ uptake.

The physiology of calcifying phytoplankton (coccolithophores) will be impacted in positive and negative ways. Photosynthesis will benefit from the higher CO₂ conditions, while calcification is projected to be negatively impacted (Bach *et al.* 2013). This mechanistic understanding of physiology derived in the laboratory can be tested in the modern Ocean (Beaufort *et al.* 2011) and fossil records (O'Dea *et al.* 2014). Globally, with increasing CO₂ in surface waters carbonate production in coccolithophores decreases (Beaufort, *et al.* 2011). A similar response can be found across the last glacial interglacial transitions but also highlights a high complexity of species specific reactions (Beaufort, *et al.* 2011; see Pörtner, *et al.* 2014 for a recent discussion).

Carbonate production in foraminifers is driven by test size and thickness. Size in foraminifers is predominantly impacted by the optimal temperature of each species (Schmidt, *et al.* 2006; Schmidt, *et al.* 2004) and therefore fluctuates with the environment through time in a predictable fashion (Schmidt *et al.* 2003). Weight, reconstructed as size normalized weight (SNW), appears to be influenced by a wider range of parameters. The carbonate produced by an individual has been related to changes in carbonate saturation in experiments (Russell *et al.* 2004) with strong species specific sensitivities in field samples (Barker and Elderfield 2002; Osborne *et al.* 2016) although many environmental parameters in the ocean are highly correlated. Seasonally, even at places without strong changes in saturation, SNW

CLIMATE CHANGE IMPACTS: THE ROLE OF PALAEOONTOLOGY

varies additionally in response to temperature and food availability again with species specific sensitivities (Weinkauf *et al.* 2016).

Such species specific responses to multiple stressors are difficult to disentangle in experiments today. Applying the fossil record of warming and acidification has the potential to provide significant advances in understanding (Osborne, *et al.* 2016) with examples focusing on the most recent glacial-interglacial cycle (Barker and Elderfield 2002) and in the Pliocene (Davis *et al.* 2013).

The *early Late Pliocene* (3.6 to 3 Ma ago) is an attractive target interval as it is the last extended interval in Earth history when atmospheric CO₂ was comparable to today (Seki *et al.* 2010) and global climate globally 3°C warmer. Pliocene SNW has shown that species react to the environmental changes associated with glacial-interglacial changes in deeper time and that their weight changes scale with the amplitude of climate change, i.e. a small Pliocene forcing results in smaller weight changes (Davis, *et al.* 2013). The shortfall of using this time interval is that it is: (1) a generally warmer world without the dramatic rate of change we currently experience; and (2) amplitudes of change were significantly smaller and therefore likely not stretching the environmental niche of the species as much.

The major strength of the fossil record though is not the assessment of individual species but the potential to combine pelagic studies with shelf species. Climate change, either due to natural decadal variability or anthropogenic change, has been shown to impact on Great Barrier Reef corals (De'ath *et al.* 2012), coralline algae (Halfar *et al.* 2011) and molluscs (Gazeau *et al.* 2013). Such skeleton-based

CLIMATE CHANGE IMPACTS: THE ROLE OF PALAEOONTOLOGY

approaches can be applied to a wide range of species and time intervals. The simplicity of the methodology provides potential for comparative work and large-scale synthesis of data from surface and deep ocean and pelagic and neritic ecosystems. A combined assessment of the impact of climate change on such a wide range of calcifiers would provide invaluable insights into vulnerabilities and potential for future carbonate production. One such target interval is the PETM, one of the Paleogene hyperthermal events ~56 myrs ago.

RAPID CLIMATE CHANGE - PALEOGENE HYPERTHERMALS

A significant body of work over the last decade or so has concentrated on assessing the impacts of transient climate change events in the geological record with the most detailed examples stemming from the Palaeogene hyperthermals. These events are often associated with large negative carbon isotope excursions, indicating the addition of vast amounts of carbon to the global carbon reservoir (Kirtland Turner and Ridgwell 2016). The injection of the carbon over short time intervals led to transient warming (Dunkley Jones, *et al.* 2013) and ocean acidification (Hönisch, *et al.* 2012). The PETM is the best studied of the hyperthermal events in a series throughout the Paleocene and Eocene. The cause for the rapid climate change leading to these transient events is still under debate (see (Frieling *et al.* 2016)); it is likely to be different for each given their different timing with regards to external forcing (Zeebe *et al.* 2017).

The power of these events, especially the PETM, is the potential for integrated assessment across latitudes and habitats from the shelf to the open ocean but also

CLIMATE CHANGE IMPACTS: THE ROLE OF PALAEOLOGY

between land and sea. The focus of the scientific endeavour to this point was on determining the physical and chemical changes associated with the climate perturbation (see Zhou *et al.* 2016; Zeebe *et al.* 2016; Gibbs *et al.* 2016 for some recent examples), though the event was first recognised by its biotic impact on the deep sea ecosystem (Thomas and Shackleton 1996).

The migration of species towards to poles and species turnover can be seen in a wide range of organisms in the marine planktic ecosystem including foraminifers (Thomas and Shackleton 1996; Lu and Keller 1995), coccolithophores (Bralower 2002) and dinoflagellates (Sluijs *et al.* 2008; Crouch *et al.* 2001) with subtropical species reaching the latitudes that are today associated with polar systems. Changes in the seasonality of runoff (McInerney and Wing 2011) and increased weathering (Dickson *et al.* 2015) led to reallocation of nutrients causing responses amongst the phytoplankton (Gibbs *et al.* 2006), including blooms of groups associated with harmful algal blooms (Sluijs and Brinkhuis 2009) but also shelf dwelling benthic foraminifers (Scheibner *et al.* 2005).

The most profound impacts were recorded in the deep sea where 40% of benthic foraminifers went extinct, a group of species that showed resistance to the mass extinction at the KPg boundary (Thomas 2007). Those benthic foraminiferal species surviving the PETM had the ability to enhance their calcification relative to body volume (Foster *et al.* 2013). Intriguingly, deep sea ostracods show highly variable responses, some of which suggest that the climate change impacted the assemblages negatively (Steineck and Thomas 1996) while others suggest little change (Webb *et al.* 2009).

CLIMATE CHANGE IMPACTS: THE ROLE OF PALAEOLOGY

The strength of this time interval is that global assessments are feasible (McInerney and Wing 2011). While the terrestrial record often does not have the same temporal resolution or accuracy of age determination, the geological record of the PETM includes high resolution sections of terrestrial sequences with astronomically tuned age models (Bowen *et al.* 2015). Those sections show rapid large plant range shifts (Wing *et al.* 2005). The mammalian records show rapid increases in mean species body size and abundance of herbivores and frugivores (Clyde and Gingerich 1998) in response to immigration of cosmopolitan new taxa (Gingerich 2003).

The PETM is an excellent example, which shows that the current response to climate change over the last decades is driven by the same underlying universal principles as in the past and demonstrates that past events are, therefore, suitable to help predict the future. Comparing the Paleogene hyperthermals and their different amplitudes of environmental change (Jennions *et al.* 2015) holds potential for assessing the importance of different amplitudes of climate change and the impact these have on both marine and terrestrial ecosystems.

OUTLOOK

Traditionally, the impacts of climate change are assessed by time-consuming taxonomic or geochemical analysis focusing on a region (Edgar *et al.* 2013) or globally within a specific time slice (Yasuhara *et al.* 2012). Rarely are different events compared and, when they are, only at a single location (Gibbs *et al.* 2012; Jennions, *et al.* 2015); this is due to the extremely time consuming work of species identification.

CLIMATE CHANGE IMPACTS: THE ROLE OF PALAEOONTOLOGY

A wealth of information could be assembled if scientists focusing on different fossil groups were to generate papers that cross traditional discipline boundaries. The power of comparing sets of events, while acknowledging our difficulties in aligning absolute dates, would provide insights into the importance of rates of change and the associated impact. Such a question, while fundamental to the size of the impact of future climate changes, is hard, and perhaps impossible, to tackle experimentally.

Selecting species with different ecologies will allow morphological changes to be linked to environmental drivers, and the sensitivities to these drivers to be tested against prior adaptation. Overcoming this regional or temporal limitation by generating global datasets covering several time slices that experienced different rates and ranges of climate change for a range of organisms has to be our goal for the next decade.

Palaeontology's contribution to the last assessment on impacts of and adaptation to climate change was highly visible in the summary for policy makers of IPCC Working Groups II (Impacts, Adaptation, and Vulnerabilities) 'Many terrestrial, freshwater, and marine species have shifted their geographic ranges, seasonal activities, migration patterns, abundances, and species interactions in response to ongoing climate change (high confidence)...While only a few recent species extinctions have been attributed as yet to climate change (high confidence), natural global climate change at rates slower than current anthropogenic climate change caused significant ecosystem shifts and species extinctions during the past millions of years (high confidence) (IPCC

2014).’ In collaborations with others we should aim for a strong visibility of our field and the knowledge it can provide in more policy relevant assessments in the future.

Acknowledgements.

DNS would like to thank the Royal Society for support in form of a Wolfson Merit Award and Jessica Pollitt for comments on this draft.

REFERENCES

- BACH, L. T., MACKINDER, L. C. M., SCHULZ, K. G., WHEELER, G., SCHROEDER, D. C., BROWNLEE, C. and RIEBESELL, U. 2013. Dissecting the impact of CO₂ and pH on the mechanisms of photosynthesis and calcification in the coccolithophore *Emiliana huxleyi*. *New Phytologist*, **199**, 121-134.
- BARKER, S. and ELDERFIELD, H. 2002. Foraminiferal calcification response to glacial-interglacial changes in atmospheric CO₂. *Science*, **297**, 833-836.
- BEAUFORT, L., PROBERT, I., DE GARIDEL-THORON, T., BENDIF, E. M., RUIZ-PINO, D., METZL, N., GOYET, C., BUCHET, N., COUPEL, P., GRELAUD, M., ROST, B., RICKABY, R. E. M. and DE VARGAS, C. 2011. Sensitivity of coccolithophores to carbonate chemistry and ocean acidification. *Nature*, **476**, 80-83.
- BEAUGRAND, G., EDWARDS, M., RAYBAUD, V., GOBERVILLE, E. and KIRBY, R. R. 2015. Future vulnerability of marine biodiversity compared with contemporary and past changes. *Nature Clim. Change*, **5**, 695-701.
- BEAUGRAND, G. and REID, P. C. 2003. Long-term changes in phytoplankton, zooplankton and salmon related to climate. *Global Change Biology*, **9**, 801-817.
- BEAUGRAND, G., REID, P. C., IBANEZ, F., LINDLEY, J. A. and EDWARDS, M. 2002. Reorganization of North Atlantic Marine Copepod Biodiversity and Climate. *Science*, **296**, 1692-1694.
- BOPP, L., RESPLANDY, L., ORR, J. C., DONEY, S. C., DUNNE, J. P., GEHLEN, M., HALLORAN, P., HEINZE, C., ILYINA, T., SÉFÉRIAN, R., TJIPUTRA, J. and VICHI, M. 2013. Multiple stressors of ocean ecosystems in the 21st century: projections with CMIP5 models. *Biogeosciences*, **10**, 6225-6245.
- BOWEN, G. J., MAIBAUER, B. J., KRAUS, M. J., ROHL, U., WESTERHOLD, T., STEIMKE, A., GINGERICH, P. D., WING, S. L. and CLYDE, W. C. 2015. Two massive, rapid releases of carbon during the onset of the Palaeocene-Eocene thermal maximum. *Nature Geosci*, **8**, 44-47.
- BRALOWER, T. J. 2002. Evidence of surface water oligotrophy during the Paleocene-Eocene thermal maximum: nannofossil assemblage data from Ocean Drilling Program Site 690, Maud Rise, Weddell Sea. *Paleoceanography*, **17**, 10.1029/2001PA000662.
- CAROMEL, A. G. M., SCHMIDT, D. N., FLETCHER, I. and RAYFIELD, E. J. 2016. Morphological change during the ontogeny of the planktic foraminifera. *Journal of Micropalaeontology*, **35**, 2-19.
- CHEUNG, W. W. L., WATSON, R. and PAULY, D. 2013. Signature of ocean warming in global fisheries catch. *Nature*, **497**, 365-368.
- CLIMAP PROJECT MEMBERS 1976. The surface of the ice-age earth. *Science*, **191**, 1131-1144.
- CLYDE, W. C. and GINGERICH, P. D. 1998. Mammalian community response to the latest Paleocene thermal maximum: An isotaphonomic study in the northern Bighorn Basin, Wyoming. *Geology*, **26**, 1011-1014.
- COLLINS, S. and BELL, G. 2004. Phenotypic consequences of 1,000 generations of selection at elevated CO₂ in a green alga. *Nature*, **431**, 566-569.
- COLLINS, S., ROST, B. and RYNEARSON, T. A. 2014. Evolutionary potential of marine phytoplankton under ocean acidification. *Evolutionary Applications*, **7**, 140-155.
- CROUCH, E. M., HEILMANN-CLAUSEN, C., BRINKHUIS, H., MORGANS, H. E. G., ROGERS, K. M., EGGER, H. and SCHMITZ, B. 2001. Global dinoflagellate event associated with the late Paleocene thermal maximum. *Geology*, **29**, 315-318.
- DAVIS, C. V., BADGER, M. P. S., BOWN, P. R. and SCHMIDT, D. N. 2013. The response of calcifying plankton to climate change in the Pliocene. *Biogeosciences*, **10**, 6131-6139.

CLIMATE CHANGE IMPACTS: THE ROLE OF PALAEOONTOLOGY

- DE'ATH, G., FABRICIUS, K. E., SWEATMAN, H. and PUOTINEN, M. 2012. The 27-year decline of coral cover on the Great Barrier Reef and its causes. *Proceedings of the National Academy of Sciences*, **109**, 17995-17999.
- DE'ATH, G., LOUGH, J. M. and FABRICIUS, K. E. 2009. Declining Coral Calcification on the Great Barrier Reef. *Science*, **323**, 116-119.
- DICKSON, A. J., COHEN, A. S., COE, A. L., DAVIES, M., SHCHERBININA, E. A. and GAVRILOV, Y. O. 2015. Evidence for weathering and volcanism during the PETM from Arctic Ocean and Peri-Tethys osmium isotope records. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **438**, 300-307.
- DOWSETT, H. J., ROBINSON, M. M., HAYWOOD, A. M., HILL, D. J., DOLAN, A. M., STOLL, D. K., CHAN, W.-L., ABE-OUCHI, A., CHANDLER, M. A., ROSENBLOOM, N. A., OTTO-BLIESNER, B. L., BRAGG, F. J., LUNT, D. J., FOLEY, K. M. and RIESSELMAN, C. R. 2012. Assessing confidence in Pliocene sea surface temperatures to evaluate predictive models. *Nature Clim. Change*, **2**, 365-371.
- DUNKLEY JONES, T., LUNT, D. J., SCHMIDT, D. N., RIDGWELL, A., SLUIJS, A., VALDES, P. J. and MASLIN, M. 2013. Climate model and proxy data constraints on ocean warming across the Paleocene-Eocene Thermal Maximum. *Earth-Science Reviews*, **125**, 123-145.
- DUPONT, S., DOREY, N., STUMPP, M., MELZNER, F. and THORNDYKE, M. 2012. Long-term and trans-life-cycle effects of exposure to ocean acidification in the green sea urchin *Strongylocentrotus droebachiensis*. *Marine Biology*, 1-9.
- DUTKIEWICZ, S., MORRIS, J. J., FOLLOWS, M. J., SCOTT, J., LEVITAN, O., DYHRMAN, S. T. and BERMAN-FRANK, I. 2015. Impact of ocean acidification on the structure of future phytoplankton communities. *Nature Clim. Change*, **5**, 1002-1006.
- EDGAR, K. M., BOHATY, S. M., GIBBS, S. J., SEXTON, P. F., NORRIS, R. D. and WILSON, P. A. 2013. Symbiont 'bleaching' in planktic foraminifera during the Middle Eocene Climatic Optimum. *Geology*, **41**, 15-18.
- EDWARDS, M., REID, P. and PLANQUE, B. 2001. Long-term and regional variability of phytoplankton biomass in the Northeast Atlantic (1960-1995). *ICES Journal of Marine Science*, **58**, 39-49.
- ENFIELD, D. B., MESTAS-NUÑEZ, A. M. and TRIMBLE, P. J. 2001. The Atlantic Multidecadal Oscillation and its relation to rainfall and river flows in the continental U.S. . *Geophysical Research Letters*, **28**, 2077-2080.
- ETHERIDGE, D. M., STEELE, L. P., LANGENFELDS, R. L., FRANCEY, R. J., BARNOLA, J. M. and MORGAN, V. I. 1996. Natural and anthropogenic changes in atmospheric CO₂ over the last 1000 years from air in Antarctic ice and firn. *Journal of Geophysical Research*, **101**, 4115-4128.
- FIELD, D. B., BAUMGARTNER, T. R., CHARLES, C. D., FERREIRA-BARTRINA, V. and OHMAN, M. D. 2006. Planktonic Foraminifera of the California Current Reflect 20th-Century Warming. *Science*, **311**, 63-66.
- FOSTER, L. C., SCHMIDT, D. N., THOMAS, E., ARNDT, S. and RIDGWELL, A. 2013. Surviving rapid climate change in the deep sea during the Paleogene hyperthermals. *Proceedings of the National Academy of Sciences*, **110**, 9273-9276.
- FRIELING, J., SVENSEN, H. H., PLANKE, S., CRAMWINCKEL, M. J., SELNES, H. and SLUIJS, A. 2016. Thermogenic methane release as a cause for the long duration of the PETM. *Proceedings of the National Academy of Sciences*, **113**, 12059-12064.
- GAZEAU, F., PARKER, L. M., COMEAU, S., GATTUSO, J.-P., O'CONNOR, W. A., MARTIN, S., PÖRTNER, H.-O. and ROSS, P. M. 2013. Impacts of ocean acidification on marine shelled molluscs. *Marine Biology*, **160**, 2207-2245.
- GIBBS, S. J., BOWN, P. R., MURPHY, B. H., SLUIJS, A., EDGAR, K. M., PÄLIKE, H., BOLTON, C. T. and ZACHOS, J. C. 2012. Scaled biotic disruption during early Eocene global warming events. *Biogeosciences*, **9**, 4679-4688.

CLIMATE CHANGE IMPACTS: THE ROLE OF PALAEOONTOLOGY

- GIBBS, S. J., BOWN, P. R., RIDGWELL, A., YOUNG, J. R., POULTON, A. J. and O'DEA, S. A. 2016. Ocean warming, not acidification, controlled coccolithophore response during past greenhouse climate change. *Geology*, **44**, 59-62.
- GIBBS, S. J., BRALOWER, T. J., BOWN, P. R., ZACHOS, J. C. and BYBELL, L. M. 2006. Shelf and open-ocean calcareous phytoplankton assemblages across the Paleocene-Eocene Thermal Maximum: Implications for global productivity gradients. *Geology*, **34**, 233-236.
- GINGERICH, P. D. 2003. Mammalian responses to climate change at the Paleocene-Eocene boundary: Polecat Bench record in the northern Bighorn Basin, Wyoming. 463-478. In WING, S. C., GINGERICH, P. D., SCHMITZ, B. and THOMAS, E. (eds). *In Causes and consequences of globally warm climates in the early Paleogene*. Geological Society of America, pp. Custom 7.
- GUISAN, A. and ZIMMERMANN, N. E. 2000. Predictive habitat distribution models in ecology. *Ecological Modelling*, **135**, 147-186.
- HALFAR, J., HETZINGER, S., ADEY, W., ZACK, T., GAMBOA, G., KUNZ, B., WILLIAMS, B. and JACOB, D. E. 2011. Coralline algal growth-increment widths archive North Atlantic climate variability. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **302**, 71-80.
- HENDERSON, G. M. 2002. New oceanic proxies for paleoclimate. *Earth and Planetary Science Letters*, **203**, 1-13.
- HENEHAN, M. J., HULL, P. M., PENMAN, D. E., RAE, J. W. B. and SCHMIDT, D. N. 2016. Biogeochemical significance of pelagic ecosystem function: An end-Cretaceous case study. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **371**.
- HENSON, S. A., BEAULIEU, C., ILYINA, T., JOHN, J. G., LONG, M., SÉFÉRIAN, R., TJIPUTRA, J. and SARMIENTO, J. L. 2017. Rapid emergence of climate change in environmental drivers of marine ecosystems. *Nature Communications*, **8**, 14682.
- HÖNISCH, B., RIDGWELL, A., SCHMIDT, D. N., THOMAS, E., GIBBS, S. J., SLUIJS, A., ZEEBE, R., KUMP, L., MARTINDALE, R. C., GREENE, S. E., KIESSLING, W., RIES, J., ZACHOS, J. C., ROYER, D. L., BARKER, S., MARCHITTO, T. M., MOYER, R., PELEJERO, C., ZIVERI, P., FOSTER, G. L. and WILLIAMS, B. 2012. The Geological Record of Ocean Acidification. *Science*, **335**, 1058-1063.
- HULL, P. M., NORRIS, R. D., BRALOWER, T. J. and SCHUETH, J. D. 2011. A role for chance in marine recovery from the end-Cretaceous extinction. *Nature Geoscience*, **4**, 856-860.
- IPCC (Ed.) 2014. *Summary for Policymakers*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 32 pp.
- JENNIONS, S. M., THOMAS, E., SCHMIDT, D. N., LUNT, D. and RIDGWELL, A. 2015. Changes in benthic ecosystems and ocean circulation in the Southeast Atlantic across Eocene Thermal Maximum 2. *Paleoceanography*, **30**, 1059-1077.
- KEELING, C. D., WHORF, T. P., WAHLEN, M. and VAN DER PLICHT, J. 1995. Interannual extremes in the rate of rise of atmospheric carbon dioxide since 1980. *Nature*, **375**, 666-670.
- KENNETT, J. P. and STOTT, L. D. 1991. Abrupt deep-sea warming, palaeoceanographic changes and benthic extinctions at the end of the Paleocene. *Nature*, **353**, 225-229.
- KIRTLAND TURNER, S. and RIDGWELL, A. 2016. Development of a novel empirical framework for interpreting geological carbon isotope excursions, with implications for the rate of carbon injection across the PETM. *Earth and Planetary Science Letters*, **435**, 1-13.
- KORTSCH, S., PRIMICERIO, R., FOSSHEIM, M., DOLGOV, A. V. and ASCHAN, M. 2015. Climate change alters the structure of arctic marine food webs due to poleward shifts of boreal generalists. *Proceedings of the Royal Society of London B: Biological Sciences*, **282**.

CLIMATE CHANGE IMPACTS: THE ROLE OF PALAEOONTOLOGY

- KROEKER, K. J., KORDAS, R. L., CRIM, R. N. and SINGH, G. G. 2010. Meta-analysis reveals negative yet variable effects of ocean acidification on marine organisms. *Ecology Letters*, **13**, 1419–1434.
- KRUEGER-HADFIELD, S. A., BALESTRERI, C., SCHROEDER, J., HIGHFIELD, A., HELAOUËT, P., ALLUM, J., MOATE, R., LOHBECK, K. T., MILLER, P. I., RIEBESELL, U., REUSCH, T. B. H., RICKABY, R. E. M., YOUNG, J., HALLEGRAEFF, G., BROWNLEE, C. and SCHROEDER, D. C. 2014. Genotyping an *Emiliana huxleyi* (prymnesiophyceae) bloom event in the North Sea reveals evidence of asexual reproduction. *Biogeosciences*, **11**, 5215–5234.
- LEFORT, S., AUMONT, O., BOPP, L., ARSOUZE, T., GEHLEN, M. and MAURY, O. 2015. Spatial and body-size dependent response of marine pelagic communities to projected global climate change. *Global Change Biology*, **21**, 154–164.
- LOHBECK, K. T., RIEBESELL, U. and REUSCH, T. B. H. 2012. Adaptive evolution of a key phytoplankton species to ocean acidification. *Nature Geoscience*, **5**, 346–351.
- LU, G. and KELLER, G. 1995. Ecological stasis and saltation: species richness change in planktic foraminifera during the late Paleocene to early Eocene, DSDP Site 577. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **117**, 211–227.
- LUNT, D. J., HAYWOOD, A. M., SCHMIDT, G. A., SALZMANN, U., VALDES, P. J. and DOWSETT, H. J. 2010a. Earth system sensitivity inferred from Pliocene modelling and data. *Nature Geoscience*, **3**, 60–64.
- LUNT, D. J., VALDES, P. J., JONES, T. D., RIDGWELL, A., HAYWOOD, A. M., SCHMIDT, D. N., MARSH, R. and MASLIN, M. 2010b. Non-linear warming of South Atlantic intermediate waters within PETM-like model simulations. *Geology*, **38**, 875–878.
- MARGO PROJECT MEMBERS 2009. Constraints on the magnitude and patterns of ocean cooling at the Last Glacial Maximum. *Nature Geoscience*, **2**, 127–132.
- MCINERNEY, F. A. and WING, S. L. 2011. The Paleocene-Eocene Thermal Maximum: A perturbation of carbon cycle, climate, and biosphere with implications for the future. *Annual Review of Earth and Planetary Sciences*, **39**, 489–516.
- MILLENNIUM ECOSYSTEM ASSESSMENT 2005. *Ecosystems and Human Well-being: General Synthesis*. World Resources Institute, Washington, DC.
- O'DEA, S. A., GIBBS, S. J., BOWN, P. R., YOUNG, J. R., POULTON, A. J., NEWSAM, C. and WILSON, P. A. 2014. Coccolithophore calcification response to past ocean acidification and climate change. *Nature Communication*, **5**.
- ORR, J. C., FABRY, V. J., AUMONT, O., BOPP, L., DONEY, S. C., FEELY, R. A., GNANADESIKAN, A., GRUBER, N., ISHIDA, A., JOOS, F., KEY, R. M., LINDSAY, K., MAIER-REIMER, E., MATEAR, R., MONFRAY, P., MOUCHET, A., NAJJAR, R. G., PLATTNER, G.-K., RODGERS, K. B., SABINE, C. L., SARMIENTO, J. L., SCHLITZER, R., SLATER, R. D., TOTTERDELL, I. J., WEIRIG, M.-F., YAMANAKA, Y. and YOOL, A. 2005. Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms. *Nature*, **437**, 681–686.
- OSBORNE, E. B., THUNELL, R. C., MARSHALL, B. J., HOLM, J. A., TAPPA, E. J., BENITEZ-NELSON, C., CAI, W.-J. and CHEN, B. 2016. Calcification of the planktonic foraminifera *Globigerina bulloides* and carbonate ion concentration: Results from the Santa Barbara Basin. *Paleoceanography*, **31**, 1083–1102.
- PEARSON, R. G. and DAWSON, T. P. 2003. Predicting the impacts of climate change on the distribution of species: are bioclimate envelope models useful? *Global Ecology and Biogeography*, **12**, 361–371.
- POLOCZANSKA, E. S., BROWN, C. J., SYDEMAN, W. J., KIESSLING, W., SCHOEMAN, D. S., MOORE, P. J., BRANDER, K., BRUNO, J. F., BUCKLEY, L. B., BURROWS, M. T., DUARTE, C. M., HALPERN, B. S., HOLDING, J., KAPPEL, C. V., O'CONNOR, M. I., PANDOLFI, J. M., PARMESAN, C., SCHWING, F., THOMPSON, S. A. and RICHARDSON, A. J. 2013. Global imprint of climate change on marine life. *Nature Clim. Change*, **3**, 919–925.

CLIMATE CHANGE IMPACTS: THE ROLE OF PALAEOONTOLOGY

- PÖRTNER, H. O., KARL, D., BOYD, P. W., CHEUNG, W., LLUCH-COTA, S. E., NOJIRI, Y., SCHMIDT, D. N. and ZAVIALOV, P. 2014. Ocean systems. 411-484. In FIELD, C. B., BARROS, V. R., DOKKEN, D. J., MACH, K. J., MASTRANDREA, M. D., BILIR, T. E., CHATTERJEE, M., EBI, K. L., ESTRADA, Y. O., GENOVA, R. C., GIRMA, B., KISSEL, E. S., LEVY, A. N., MACCRACKEN, S., MASTRANDREA, P. R. and WHITE, L. L. (eds). *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 6 pp. Custom 7.
- RIDGWELL, A. and SCHMIDT, D. N. 2010. Past constraints on the vulnerability of marine calcifiers to massive carbon dioxide release. *Nature Geoscience*, **3**, 196-200.
- ROBINSON, M. M., DOWSETT, H. J., DWYER, G. S. and LAWRENCE, K. T. 2008. Reevaluation of mid-Pliocene North Atlantic sea surface temperatures. *Paleoceanography*, **23**, PA3213, <http://dx.doi.org/10.1029/2008PA001608>.
- ROY, T., LOMBARD, F., BOPP, L. and GEHLEN, M. 2015. Projected impacts of climate change and ocean acidification on the global biogeography of planktonic Foraminifera. *Biogeosciences*, **12**, 2873-2889.
- RUSSELL, A. D., HÖNISCH, B., SPERO, H. J. and LEA, D. W. 2004. Effects of seawater carbonate ion concentration and temperature on shell U, Mg, and Sr in cultured planktonic foraminifera. *Geochimica et Cosmochimica Acta*, **68**, 4347-4361.
- SCHEIBNER, C., SPEIJER, R. P. and MARZOUK, A. M. 2005. Turnover of larger foraminifera during the Paleocene-Eocene Thermal Maximum and paleoclimatic control on the evolution of platform ecosystems. *Geology*, **33**, 493-496.
- SCHEININ, M., RIEBESELL, U., RYNEARSON, T. A., LOHBECK, K. T. and COLLINS, S. 2015. Experimental evolution gone wild. *Journal of the Royal Society Interface*, **12**.
- SCHIEBEL, R. 2002. Planktic foraminiferal sedimentation and the marine calcite budget. *Global biogeochemical cycles*, **16**, doi: 10.1029/2001GB001459.
- SCHMIDT, D. N., LAZARUS, D., YOUNG, J. and KUCERA, M. 2006. Biogeography and evolution of body-size of marine plankton. *Earth-Science Reviews*, **78**, 239-266.
- SCHMIDT, D. N., RENAUD, S. and BOLLMANN, J. 2003. Response of planktic foraminiferal size to late Quaternary climate change. *Paleoceanography*, **18**, 10.1029/2002PA000831.
- SCHMIDT, D. N., RENAUD, S., BOLLMANN, J., SCHIEBEL, R. and THIERSTEIN, H. R. 2004. Size distribution of Holocene planktic foraminifer assemblages: biogeography, ecology and adaptation. *Marine Micropaleontology*, **50**, 319-338.
- SEKI, O., FOSTER, G. L., SCHMIDT, D. N., MACKENSEN, A., KAWAMURA, K. and PANCOST, R. D. 2010. Alkenone and boron-based Pliocene pCO₂ records. *Earth and Planetary Science Letters*, **292**, 201-211.
- SLUIJS, A. and BRINKHUIS, H. 2009. A dynamic climate and ecosystem state during the Paleocene-Eocene Thermal Maximum: inferences from dinoflagellate cyst assemblages on the New Jersey Shelf. *Biogeosciences*, **6**, 1755-1781.
- SLUIJS, A., ROEHL, U., SCHOUTEN, S., BRUMSACK, H.-J., SANGIORGI, F., SINNINGHE DAMSTE, J. S. and BRINKHUIS, H. 2008. Arctic late Paleocene-early Eocene paleoenvironments with special emphasis on the Paleocene-Eocene thermal maximum (Lomonosov Ridge, Integrated Ocean Drilling Program Expedition 302). *Paleoceanography*, **23**, PA1S11.
- STANTON, J. C., SHOEMAKER, K. T., PEARSON, R. G. and AKÇAKAYA, H. R. 2014. Warning times for species extinctions due to climate change. *Global Change Biology*, n/a-n/a.
- STEINECK, P. L. and THOMAS, E. 1996. The latest Paleocene crisis in the deep sea: Ostracode succession at Maud Rise, Southern Ocean. *Geology*, **24**, 583-586.

CLIMATE CHANGE IMPACTS: THE ROLE OF PALAEONTOLOGY

- STOCKER, T. F., QIN, D., PLATTNER, G.-K., TIGNOR, M., ALLEN, S. K., BOSCHUNG, J., NAUELS, A., XIA, Y., BEX, V. and MIDGLEY, P. M. 2013. Working group I contribution to the IPCC 5th assessment report. *Climate change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, United Kingdom.
- THOMAS, E. 2003. Extinction and food at the seafloor: A high-resolution benthic foraminiferal record across the Initial Eocene Thermal Maximum, Southern Ocean Site 690. 319-332. In WING, S. L., GINGERICH, P. D., SCHMITZ, B. and THOMAS, E. (eds). *Causes and Consequences of Globally Warm Climates in the Early Paleogene. Geological Society of America Special Paper 369*. Geological Society of America, Boulder, Colorado, pp. Custom 7.
- 2007. Cenozoic mass extinctions in the deep sea: what perturbs the largest habitat on earth? 1-23. In MONECHI, S., COCCIONI, R. and RAMPINO, M. R. (eds). *Geological Society of America Special Paper*. Geological Society of America, pp. Custom 7.
- THOMAS, E. and SHACKLETON, N. J. 1996. The Palaeocene-Eocene benthic foraminiferal extinction and stable isotope anomalies. 401-441. In KNOX, R. W. O. B., CORFIELD, R. M. and DUNAY, R. E. (eds). *Correlation of the Early Paleogene in Northwestern Europe, Geological Society London Special Publication, 101*. Geological Society of London, London, United Kingdom, pp. Custom 7.
- THOMSEN, J., GUTOWSKA, M. A., SAPHÖRSTER, J., HEINEMANN, A., TRÜBENBACH, K., FIETZKE, J., HIEBENTHAL, C., EISENHAUER, A., KÖRTZINGER, A., WAHL, M. and MELZNER, F. 2010. Calcifying invertebrates succeed in a naturally CO₂-rich coastal habitat but are threatened by high levels of future acidification. *Biogeosciences*, **7**, 3879-3891.
- TINDALL, J., FLECKER, R., VALDES, P., SCHMIDT, D. N., MARKWICK, P. and HARRIS, J. 2010. Modeling the oxygen isotope distribution of ancient seawater using a coupled ocean-atmosphere GCM: implications for reconstructing early Eocene climate. *Earth and Planetary Science Letters*, **292**, 265-273.
- TURLEY, C., EBY, M., RIDGWELL, A. J., SCHMIDT, D. N., BROWNLEE, C., FINDLAY, H. S., FABRY, V. J., FEELY, R. A., RIEBESELL, U. and GATTUSO, J.-P. 2010. The Societal challenge of ocean acidification. *Marine Pollution Bulletin*, **60**, 787-792.
- WALL, M., RAGAZZOLA, F., FOSTER, L. C., FORM, A. and SCHMIDT, D. N. 2015. pH up-regulation as a potential mechanism for the cold-water coral *Lophelia pertusa* to sustain growth in aragonite undersaturated conditions. *Biogeosciences*, **12**, 6869-6880.
- WALTHER, G.-R., POST, E., CONVEY, P., MENZEL, A., PARMESAN, C., BEEBEE, T. J. C., FROMENTIN, J.-M., HOEGH-GULDBERG, O. and BAIRLEIN, F. 2002. Ecological responses to recent climate change. *Nature*, **416**, 389-395.
- WATERSON, A. M., EDGAR, K. M., SCHMIDT, D. N. and VALDES, P. J. 2017. Quantifying the stability of planktic foraminiferal physical niches between the Holocene and Last Glacial Maximum. *Paleoceanography*, **32**, 74-89.
- WATERSON, A. M., SCHMIDT, D. N., VALDES, P. J., HOLROYD, P. A., NICHOLSON, D. B., FARNSWORTH, A. and BARRETT, P. M. 2016. Modelling the testudine climatic niche: a deep-time perspective *Proceedings of the Royal Society B*, **283**, 20161408.
- WEBB, A. E., LEIGHTON, L. R., SCHELLENBERG, S. A., LANDAU, E. A. and THOMAS, E. 2009. Impact of the Paleocene-Eocene thermal maximum on deep-ocean microbenthic community structure: Using rank-abundance curves to quantify paleoecological response. *Geology*, **37**, 783-786.
- WEINKAUF, M. F. G., KUNZE, J. G., WANIEK, J. J. and KUČERA, M. 2016. Seasonal Variation in Shell Calcification of Planktonic Foraminifera in the NE Atlantic Reveals

CLIMATE CHANGE IMPACTS: THE ROLE OF PALAEOONTOLOGY

- Species-Specific Response to Temperature, Productivity, and Optimum Growth Conditions. *PLOS ONE*, **11**, e0148363.
- WILSON, R., TUDHOPE, A., BROHAN, P., BRIFFA, K., OSBORN, T. and TETT, S. 2006. Two-hundred-fifty years of reconstructed and modeled tropical temperatures. *Journal of Geophysical Research - Oceans* **111**, C10007.
- WING, S. L., HARRINGTON, G. J., SMITH, F. A., BLOCH, J. I., BOYER, D. M. and FREEMAN, K. H. 2005. Transient Floral Change and Rapid Global Warming at the Paleocene-Eocene Boundary. *Science*, **310**, 993-996.
- WOOD, H. L., SPICER, J. I. and WIDDICOMBE, S. 2008. Ocean acidification may increase calcification rates, but at a cost. *Proceedings of the Royal Society B: Biological Sciences*, **275**, 1767-1773.
- YASUHARA, M., HUNT, G., DOWSETT, H. J., ROBINSON, M. M. and STOLL, D. K. 2012. Latitudinal species diversity gradient of marine zooplankton for the last three million years. *Ecology Letters*, **15**, 1174-1179.
- ZEEBE, R. E., RIDGWELL, A. and ZACHOS, J. C. 2016. Anthropogenic carbon release rate unprecedented during the past 66 million years. *Nature Geoscience*, **9**, 325-329.
- ZEEBE, R. E., WESTERHOLD, T., LITTLER, K. and ZACHOS, J. C. 2017. Orbital forcing of the Paleocene and Eocene carbon cycle. *Paleoceanography*, **32**, 440–465.
- ZEEBE, R. E., ZACHOS, J. C. and DICKENS, G. R. 2009. Carbon dioxide forcing alone insufficient to explain Palaeocene-Eocene Thermal Maximum warming. *Nature Geoscience*, **2**, 576-580.
- ZHOU, X., THOMAS, E., WINGUTH, A. M. E., RIDGWELL, A., SCHER, H., HOOGAKKER, B. A. A., RICKABY, R. E. M. and LU, Z. 2016. Expanded oxygen minimum zones during the late Paleocene-early Eocene: Hints from multiproxy comparison and ocean modeling. *Paleoceanography*, **31**, 1532–1546.




Figure 1: Developmental history of 4 species of planktic foraminifera reconstructed using tomography (Caromel *et al.* 2016) and unpubl. Data. From left to right: modern, Eocene and Paleocene specimen. Each colour represents a new ontogenetic stage.

Figure 2: Environmental changes (top) and associated biological responses (bottom) for the Paleocene–Eocene Thermal Maximum (PETM, left) and the industrial era (right) from IPCC AR5 WGII, Chapter 6 (Pörtner, *et al.* 2014). Episodes of largest environmental change are indicated with darker bands. Note the different timescale between the two columns. Both time intervals are characterized by rapid warming both on land and in the ocean: {modern: Wilson *et al.* (2006) and PETM: Kennett and Stott (1991)) and increases in CO₂ (modern: (Etheridge *et al.* 1996; Keeling *et al.* 1995) and PETM: (Zeebe *et al.* 2009; LOSCAR model); (Ridgwell and Schmidt 2010; Grid Enabled Integrated Earth System Model, GENIE model). For the recent industrial era, the Atlantic Multi-decadal Oscillation (AMO) is shown to highlight an example of high-frequency sea surface temperature fluctuations (Enfield *et al.* 2001) and their influence on marine biota. Note the species-specific calcification responses to climate change with decreases, increases, and high variability (coralline alga: (Halfar, *et al.* 2011); coral: (De'ath *et al.* 2009); PETM: (Foster, *et al.* 2013). While there was extinction

CLIMATE CHANGE IMPACTS: THE ROLE OF PALAEOONTOLOGY

during the PETM (Thomas 2003), there is currently no evidence for climate-related extinction in the marine record. Warming led to migration of warm-water species into previous cold-water habitats (modern: (Field, et al. 2006); PETM: (Bralower 2002)). Pictures are examples of organisms highlighting the processes in each panel, and are not to scale.